

Agricultural Nutrient Management Program
Department of Environmental Science and Technology
0116 Symons Hall
7998 Regents Dr.
College Park, MD 20742
TEL 301-405-1319 | FAX 301-314-7375
www.extension.umd.edu/anmp

Ammonium Conservation Coefficients

Background:

Previous UME guidance recognized that tillage could be used after manure incorporation to conserve ammonium-N by limiting ammonia volatilization. However, there was no difference between various degrees of tillage intensity. Since quantitative data are lacking to assign conservation factors to specific types of tillage, we decided to use residue cover as a surrogate for incorporation and therefore amount of ammonia conservation achieved.

Tillage Information:

Conventional tillage:

- For the purpose of nutrient management planning, this will include any tillage that leaves <30% residue cover
- Examples include chisel-disc-chisel or LandsmanTM
- ammonium conservation maximized

Conservation tillage:

- Includes any tillage that leaves between 30 and 70% surface residue
- Assumed to be approximately 25 75% as effective as conventional tillage at conserving ammonium-N
- Mid-range value of 50% used to estimate conservation factor
- Examples include light discing, plug or spike aerators (such as AerwayTM), seed bed conditioners and vertical till (such as TurbotillTM)

No-till:

- Ammonium conservation minimized
- Residue cover >70%

Ammonia conservation factors for liquid manures and organic residuals (<10% dry matter, >90% moisture)

Time to	Conventional	Conservation	No-till or
incorporation	tillage	tillage	tillage > 3
			days
inject	1.0	N/A	
< 1 hr	.95	.70	
1-3	.90	.68	
3-6	.75	.60	
6-12	.65	.55	
12-24	.60	.53	
1-2 days	.55	.50	
2-3 days	.50	.48	
>3 days (no-till)			.45

Ammonia conservation factors for solid manures and organic residuals (>10% dry matter, <90% moisture)

Time to	Conventional	Conservation	No till or
incorporation	tillage	tillage	tillage > 3
			days
< 1 hr	.96	.66	
1-3	.93	.64	
3-6	.78	.57	
6-12	.71	.53	
12-24	.63	.49	
1-2 days	.58	.47	
2-3 days	.53	.44	
>3 days (no-till)			.35

Ammonia conservation factors for poultry litter

Time to	Conventional	Conservation	No till or
incorporation	tillage	tillage	tillage >14
			days
< 1 day	.97	.85	
1-2 days	.92	.82	
3 days	.88	.80	
4 days	.84	.78	
5 days	.81	.77	
6 or 7 days	.77	.75	
8 – 14 days	.74	.73	
>14 days (no-till)			.72